

Independence of the geometric Galois action on ℓ -adic cohomologies when ℓ varies

G. Böckle, W. Gajda and S. Petersen

February 28, 2013

Abstract

Let k be an algebraically closed field of arbitrary characteristic, let K/k be a finitely generated field extension and let X be a separated scheme of finite type over K . For each prime ℓ , the absolute Galois group of K acts on the ℓ -adic étale cohomology modules of X . We prove that this family of representations varying over ℓ is almost independent in the sense of Serre, i.e., that the fixed fields inside an algebraic closure of K of the kernels of the representations for all ℓ become linearly disjoint over a finite extension of K . In doing this, we also prove a number of interesting facts on the images and ramification of this family of representations.

1 Introduction

Let G be a profinite group and L a set of prime numbers. For every $\ell \in L$ let G_ℓ be a profinite group and $\rho_\ell : G \rightarrow G_\ell$ a homomorphism. Denote by

$$\rho : G \rightarrow \prod_{\ell \in L} G_\ell$$

the homomorphism induced by the ρ_ℓ . Following the notation in [35] we call the family $(\rho_\ell)_{\ell \in L}$ *independent* if $\rho(G) = \prod_{\ell \in L} \rho_\ell(G)$. The family $(\rho_\ell)_{\ell \in L}$ is said to be *almost independent* if there exists an open subgroup H of G such that $\rho(H) = \prod_{\ell \in L} \rho_\ell(H)$.

The main examples of such families of homomorphisms arise as follows: Let K be a field with algebraic closure $\tilde{K}$ and absolute Galois group $\text{Gal}(K) = \text{Aut}(\tilde{K}/K)$. Let X/K be a separated algebraic scheme¹ and denote by $\mathbb{L}$ the set of all prime numbers. For every $q \in \mathbb{N}$ and every $\ell \in \mathbb{L} \setminus \{\text{char}(k)\}$ we shall consider the representations

$$\rho_{\ell,X}^{(q)} : \text{Gal}(K) \rightarrow \text{Aut}_{\mathbb{Q}_\ell}(H^q(X_{\tilde{K}}, \mathbb{Q}_\ell)) \quad \text{and} \quad \rho_{\ell,X,c}^{(q)} : \text{Gal}(K) \rightarrow \text{Aut}_{\mathbb{Q}_\ell}(H_c^q(X_{\tilde{K}}, \mathbb{Q}_\ell))$$

of $\text{Gal}(K)$ on the étale cohomology groups $H^q(X_{\tilde{K}}, \mathbb{Q}_\ell)$ and $H_c^q(X_{\tilde{K}}, \mathbb{Q}_\ell)$. The following independence result has recently been obtained.

⁰**2010 MSC:** 11G10, 14F20.

⁰**Key words:** Galois representation, étale cohomology, algebraic scheme, finitely generated field.

¹A scheme X/K is algebraic if the structure morphism $X \rightarrow \text{Spec } K$ is of finite type (cf. [15, Def. 6.4.1]).

Theorem 1.1. *Let K be a finitely generated extension of $\mathbb{Q}$ and X/K a separated algebraic scheme. Then the families $(\rho_{\ell,X}^{(q)})_{\ell \in \mathbb{L}}$ and $(\rho_{\ell,X,c}^{(q)})_{\ell \in \mathbb{L}}$ are almost independent.*

The proof of this statement in the important special case $\text{trdeg}(K/\mathbb{Q}) = 0$ is due to Serre (cf. [35]). The case $\text{trdeg}(K/\mathbb{Q}) > 0$ was worked out in [14], answering a question of Serre (cf. [35], [36]) and Illusie [21].

The present article is concerned with a natural variant of Theorem 1.1 that grew out of the study of independence of families over fields of positive characteristic. For K a finitely generated extension of $\mathbb{F}_p$ it has long been known, e.g. [20] or [11], that the direct analogue of Theorem 1.1 is false: If $\varepsilon_\ell : \text{Gal}(\mathbb{F}_p) \rightarrow \mathbb{Z}_\ell^\times$ denotes the ℓ -adic cyclotomic character that describes the Galois action on ℓ -power roots of unity, then it is elementary to see that the family $(\varepsilon_\ell)_{\ell \in \mathbb{L} \setminus \{p\}}$ is not almost independent. It follows from this that for every abelian variety A/K , if we denote by $\sigma_{\ell,A} : \text{Gal}(K) \rightarrow \text{Aut}_{\mathbb{Q}_\ell}(T_\ell(A))$ the representation of $\text{Gal}(K)$ on the ℓ -adic Tate module of A , then $(\sigma_{\ell,A})_{\ell \in \mathbb{L} \setminus \{p\}}$ is *not* almost independent. One is thus led to study independence over the compositum $\tilde{\mathbb{F}}_p K$ obtained from the field K by adjoining all roots of unity. Having gone that far, it is then natural to study independence over any field K that is finitely generated over an arbitrary algebraically closed field k . Our main result is the following independence theorem.

Theorem 1.2. (cf. Theorem 7.5) *Let k be an algebraically closed field of characteristic $p \geq 0$. Let K/k be a finitely generated extension. Let X/K be a separated algebraic scheme. Then the families $(\rho_{\ell,X}^{(q)}|_{\text{Gal}(K)})_{\ell \in \mathbb{L} \setminus \{p\}}$ and $(\rho_{\ell,X,c}^{(q)}|_{\text{Gal}(K)})_{\ell \in \mathbb{L} \setminus \{p\}}$ are almost independent.*

It will be clear that many techniques of the present article rely on [35]. Also, some of the key results of [14] will be important. The new methods in comparison with the previous results are the following: (i) The analysis of the target of our Galois representations, reductive algebraic groups over $\mathbb{Q}_\ell$, will be based on a structural result by Larsen and Pink (cf. [27]) and no longer as for instance in [35] on extensions of results by Nori (cf. [31]). This facilitates greatly the passage from $\text{Gal}(K)$ to $\text{Gal}(K\tilde{k})$ when studying their image under $\rho_{\ell,X,?}^{(q)}$. (ii) Since we also deal with cases of positive characteristic, ramification properties will play a crucial role to obtain necessary finiteness properties of fundamental groups. The results on alterations by de Jong (cf. [6]) will obviously be needed. However we were unable to deduce all needed results from there, despite some known semistability results that follow from [6]. Instead we carry out a reduction to the case where K is absolutely finitely generated and where X/K is smooth and projective (this uses again [6]). (iii) In the latter case, we use a result by Kerz-Schmidt-Wiesend (cf. [24]) that allows one to control ramification on X by controlling it on all smooth curves on X . Since X is smooth, results of Deligne show that the semisimplifications of $\rho_{\ell,X,?}^{(q)}$ form a pure and strictly compatible system. On curves, we can then apply the global Langlands correspondence proved by Lafforgue ([26]) to obtain the required ramification properties of $(\rho_{\ell,X,?}^{(q)})_{\ell \in \mathbb{L} \setminus \{p\}}$.

Part (i) is carried out in Section 3. Results on fundamental groups and first results on ramification are the theme of Section 4; here parts of (ii) are carried out and we also refine some results from [24]). Section 5 provides the basic independence criterion on which our proof of Theorem 1.2 ultimately rests. Section 6 performs the reductions mentioned in (ii). The ideas described in (iii) are concluded in Section 7, where a slightly more precise form of Theorem 1.2 is proved.

We would like to point out that an alternative method for the part (ii) of our approach could be based on recent unpublished result by Orgogozo which proves a global semistable reduction theorem (cf. [32, 2.5.8. Prop.]). In February 2013 when our paper was complete we were informed by Anna Cadoret that together with Akio Tamagawa she has proven our Theorem 1.2 by a different method cf. [5].

Acknowledgments: G.B. thanks the Fields Institute for a research stay in the spring of 2012 during which part of this work was written. He also thanks Adam Mickiewicz University in Poznań for making possible a joint visit of the three authors in the fall of 2012. He is supported by a grant of the DFG within the SPP 1489. W.G. thanks Interdisciplinary Center for Scientific Computing (IWR) at Heidelberg University for hospitality during a research visit in January 2012 shortly after this project had been started. He was partially supported by the Alexander von Humboldt Foundation and a research grant of the National Centre of Sciences of Poland. S.P. thanks Mathematics Department at Adam Mickiewicz University for hospitality and support during several research visits.

2 Notation

For a field K with algebraic closure $\tilde{K}$, we denote by $K_s \subset \tilde{K}$ a separable closure. Then $\text{Gal}(K)$ is equivalently defined as $\text{Gal}(K_s/K)$ and as $\text{Aut}(\tilde{K}/K)$, since any field automorphism of K_s fixing K has a unique extension to $\tilde{K}$. If E/K is an arbitrary field extension, and if $\tilde{K}$ is chosen inside $\tilde{E}$, then there is a natural isomorphism $\text{Aut}(\tilde{K}/\tilde{K} \cap E) \xrightarrow{\sim} \text{Aut}(\tilde{K}E/E)$. Composing its inverse with the natural restriction $\text{Gal}(E) \rightarrow \text{Aut}(\tilde{K}E/E)$ one obtains a canonical map which we denote $\text{res}_{E/K} : \text{Gal}(E) \rightarrow \text{Gal}(K)$. If E/K is algebraic, then $\text{res}_{E/K}$ is injective and we often identify $\text{Gal}(E)$ with the subgroup $\text{res}_{E/K}(\text{Gal}(E)) = \text{Gal}(E \cap \tilde{K})$ of $\text{Gal}(K)$.

Let G be a profinite group. A *normal series* in G is a sequence

$$G \triangleright N_1 \triangleright N_2 \triangleright \cdots \triangleright N_s = \{e\}$$

of closed subgroups such that each N_i is normal in G .

A K -variety X is a scheme X that is integral separated and algebraic over K . Let S be a normal connected scheme with function field K . A separable algebraic extension E/K is said to be *unramified along S* if for every finite extension F/K inside E the normalization of S in F is étale over S . We usually consider S as a scheme equipped with the generic geometric base point $s : \text{Spec}(\tilde{K}) \rightarrow S$ and denote by $\pi_1(S) := \pi_1(S, s)$ the étale fundamental group of S . If Ω denotes the maximal extension of K in K_s which is unramified along S , then $\pi_1(S)$ can be identified with the Galois group $\text{Gal}(\Omega/K)$. A homomorphism $\rho : \text{Gal}(K) \rightarrow H$ is said to be *unramified along S* if the fixed field $K_s^{\ker(\rho)}$ is unramified along S . If E/K is an arbitrary algebraic extension, then $\rho|_{\text{Gal}(E)}$ stands for $\rho \circ \text{res}_{E/K}$.

3 Concepts from group theory

In this section, we prove a structural results of compact profinite subgroups of linear algebraic groups over $\mathbb{Q}_\ell$ (cf. Theorem 3.4) that will be crucial for the proof of the main theorem of this article. It is a consequence of a variant (cf. Proposition 3.8) of a theorem of Larsen and Pink (cf. [27, Thm. 0.2, p. 1106]). The proof of Proposition 3.8 makes strong use of the results and methods in [27], and in particular does not depend on the classification of finite simple groups.

Definition 3.1. For $c \in \mathbb{N}$ we denote by $\Sigma_\ell(c)$ the class of profinite groups M which possess a normal series by open subgroups

$$M \triangleright I \triangleright P \triangleright \{1\}$$

such that M/I is a finite product of finite simple groups of Lie type in characteristic ℓ , the group I/P is finite abelian of order prime to ℓ and index $[I : P] \leq c$, and P is a pro- ℓ group.

Definition 3.2. For $d \in \mathbb{N}$ and ℓ a prime we denote by $\text{Jor}_\ell(d)$ the class of finite groups H which possess a normal abelian subgroup N of order prime to ℓ and of index $[H : N] \leq d$. We define $\text{Jor}(d)$ as the union of the $\text{Jor}_\ell(d)$ over all primes ℓ .

Definition 3.3. A profinite group G is called n -bounded at ℓ if there exist closed compact subgroups $G_1 \subset G_2 \subset \text{GL}_n(\mathbb{Q}_\ell)$ such that G_1 is normal in G_2 and $G \cong G_1/G_2$.

The following is the main result of this section.

Theorem 3.4. For every $n \in \mathbb{N}$ there exists a constant $J'(n)$ (independent of ℓ) such that the following holds: For any prime ℓ , any group G that is n -bounded at ℓ lies in a short exact sequence

$$1 \rightarrow M \rightarrow G \rightarrow H \rightarrow 1$$

such that M is open normal in G and lies in $\Sigma_\ell(2^n)$ and H lies in $\text{Jor}_\ell(J'(n))$.

We state an immediate corollary:

Corollary 3.5. Let G be n -bounded at ℓ and define G_ℓ^+ as the normal hull of all pro- ℓ Sylow subgroups of G . Then for $\ell > J'(n)$, the group G_ℓ^+ is an open normal subgroup of M of index at most 2^n .

In the remainder of this section we shall give a proof of Theorem 3.4. Moreover we shall derive some elementary permanence properties for the property described by $\Sigma_\ell(d)$ and $\text{Jor}_\ell(d)$.

The content of the following lemma is presumably well-known.

Lemma 3.6. For every $r \in \mathbb{N}$, every algebraically closed field F and every semisimple algebraic group G of rank r the center Z of G satisfies $|Z(F)| \leq 2^r$.

Proof. Lacking a precise reference, we include a proof for the reader's convenience. Observe first that the center Z is a finite (cf. [30, I.6.20, p. 43]) diagonalizable algebraic group. Let T be a maximal torus of G . Denote by $X(T) = \text{Hom}(T, \mathbb{G}_m)$ the character group of T and by $\Phi \subset X(T)$ the set of roots of G . Then $\mathcal{R} = (X(T) \otimes \mathbb{R}, \Phi)$ is a root system. Let $P = \mathbb{Z}\Phi$ be the root lattice and Q the weight lattice of this root system. Then $P \subset X(T) \subset Q$. The center Z of G is the kernel of the adjoint representation (cf. [30, I.7.12, p. 49]). Hence $Z = \bigcap_{\chi \in \Phi} \ker(\chi)$ and there is an exact sequence

$$0 \rightarrow Z \rightarrow T \rightarrow \prod_{\chi \in \Phi} \mathbb{G}_m$$

where the right hand map is induced by the characters $\chi : T \rightarrow \mathbb{G}_m$ ($\chi \in \Phi$). We apply the functor $\text{Hom}(-, \mathbb{G}_m)$ and obtain an exact sequence

$$\prod_{\chi \in \Phi} \mathbb{Z} \rightarrow X(T) \rightarrow \text{Hom}(Z, \mathbb{G}_m) \rightarrow 0$$

The cokernel of the left hand map is $X(T)/P$. Thus $|Z(F)| \leq [X(T) : P] \leq [Q : P]$.

Furthermore, the root system $\mathcal{R}$ decomposes into a direct sum

$$\mathcal{R} = \bigoplus_{i=1}^s (E_i, \Phi_i)$$

of indecomposable root systems $\mathcal{R}_i := (E_i, \Phi_i)$. Let $r_i = \dim(E_i)$ be the rank of $\mathcal{R}_i$. Let P_i be the root lattice and Q_i the weight lattice of $\mathcal{R}_i$. Note that by definition $P = \bigoplus_i P_i$ and $Q = \bigoplus_i Q_i$. It follows from the classification of indecomposable root systems that $|Q_i/P_i| \leq 2^{r_i}$ (cf. [30, Table 9.2, p. 72]) for all i . Hence $|Z(F)| \leq |Q/P| \leq 2^{r_1} 2^{r_2} \dots 2^{r_s} = 2^r$ as desired. $\square$

Remark 3.7. *The semisimple algebraic group $(\text{SL}_{2,\mathbb{C}})^r$ has rank r and its center $(\mu_2)^r$ has exactly 2^r $\mathbb{C}$ -rational points. Hence the bound of Lemma 3.6 cannot be improved.*

The following result is an adaption of the main result of [27] by Larsen and Pink.

Proposition 3.8. *For every $n \in \mathbb{N}$, there exists a constant $J'(n)$ such that for every field F of positive characteristic ℓ and every finite subgroup Γ of $GL_n(F)$, there exists a normal series*

$$\Gamma \triangleright L \triangleright M \triangleright I \triangleright P \triangleright \{1\}$$

of Γ with the following properties:

- i) $[\Gamma : L] \leq J'(n)$.
- ii) The group L/M is abelian of order prime to ℓ .
- iii) The group M/I is a finite product of finite simple groups of Lie type in characteristic ℓ .
- iv) The group I/P is abelian of order prime to ℓ and $[I : P] \leq 2^n$.

v) P is an ℓ -group.

Furthermore the constant $J'(n)$ is the same as in [27, Thm. 0.2, p. 1106].

Proof. We can assume that F is algebraically closed. Let $J'(n)$ be the constant from [27, Thm. 0.2, p. 1106]. Larsen and Pink construct in the proof of their Theorem [27, Thm. 0.2, p. 1155–1156] a smooth algebraic group G over F containing Γ and normal subgroups Γ_i of Γ such that there is a normal series

$$\Gamma \triangleright \Gamma_1 \triangleright \Gamma_2 \triangleright \Gamma_3 \triangleright \{1\}$$

and such that $[\Gamma : \Gamma_1] \leq J'(n)$, Γ_1/Γ_2 is a product of finite simple groups of Lie type in characteristic ℓ , Γ_2/Γ_3 is abelian of order prime to ℓ and Γ_3 is an ℓ -group. Let R be the unipotent radical of the connected component G° of G . The proof of Larsen and Pink shows that $\Gamma_1 \triangleleft G^\circ(F)$, $\Gamma_3 = \Gamma \cap R(F)$ and Γ_2/Γ_3 is contained in $\overline{Z}(F)$ where $\overline{Z}$ denotes the center of the reductive group $\overline{G} := G^\circ/R$. Let $\overline{D} = [\overline{G}, \overline{G}]$ be the derived group of $\overline{G}$ and $D = [G^\circ, G^\circ]R$.

Now define $L = \Gamma_1$, $M = \Gamma_1 \cap D(F)$, $I = \Gamma_2 \cap D(F)$ and $P = \Gamma_3$. These groups are normal in Γ , because $D(F)$ is characteristic in $G^\circ(F)$ and because $\Gamma_1, \Gamma_2, \Gamma_3$ are normal in Γ . The group L/M is a subgroup of the abelian group $G^\circ(F)/D(F)$. The group M/I is a normal subgroup of Γ_1/Γ_2 , hence it is a product of finite simple groups of Lie type in characteristic ℓ . The group I/P is a subgroup of Γ_2/Γ_3 , hence I/P is abelian of order prime to ℓ . Furthermore $I/P = I/\Gamma_3$ is a subgroup of $\overline{G}(F)$ which lies in $\overline{D}(F)$ and in $\overline{Z}(F)$. Thus I/P lies in the center $\overline{Z}(F) \cap \overline{D}(F)$ of the semisimple group $\overline{D}(F)$. It follows by Lemma 3.6 that $[I : P] \leq 2^{\text{rk}(\overline{D})}$.

It remains to show that $\text{rk}(\overline{D}) \leq n$. Let T be a maximal torus of $\overline{D}$ and denote by $\pi : G^\circ \rightarrow \overline{G}$ the canonical projection. Then the algebraic group $B := \pi^{-1}(T)$ sits in an exact sequence

$$0 \rightarrow R \rightarrow B \rightarrow T \rightarrow 0$$

and B is connected smooth and solvable, because R and T have these properties. The above exact sequence splits (cf. [10, XVII.5.1]); hence B contains a copy of T . This copy is contained in a maximal torus T' of $\text{GL}_{n,F}$. Thus $n = \dim(T') \geq \dim(T) = \text{rk}(\overline{D})$ as desired. $\square$

Proof of Theorem 3.4. Suppose G is n -bounded at ℓ , so that it is a quotient G_2/G_1 with $G_i \subset \text{GL}_n(\tilde{\mathbb{Q}}_\ell)$. By Lemma 3.9(a) below, it will suffice to prove the theorem for G_2 . Thus we assume that G is a compact profinite subgroup of $\text{GL}_n(\tilde{\mathbb{Q}}_\ell)$. By compactness of G and a Baire category type argument (cf. [12, proof of Cor. 5]) the group G is contained in $\text{GL}_n(E)$ for some finite extension E of $\mathbb{Q}_\ell$. Let $\mathcal{O}_E$ be the ring of integers of the local field E . Again by compactness of G one can then find an $\mathcal{O}_E$ -lattice in E^n that is stable under G . Hence we may assume that G is a closed subgroup of $\text{GL}_n(\mathcal{O}_E)$.

Let $\mathfrak{p}$ be the maximal ideal of the local ring $\mathcal{O}_E$ and let $\mathbb{F} = \mathcal{O}_E/\mathfrak{p}$ be its residue field. The kernel $\mathcal{K}$ of the canonical map $p : \text{GL}_n(\mathcal{O}_E) \rightarrow \text{GL}_n(\mathbb{F})$ is a pro- ℓ group. Hence $Q = \mathcal{K} \cap G$ is pro- ℓ and open normal in G . We now apply Proposition 3.8 to the subgroup G/Q of $\text{GL}_n(\mathbb{F}) \subset \text{GL}_n(F)$ with $F = \overline{\mathbb{F}} \cong \overline{\mathbb{F}}_\ell$. This yields a normal series

$$G \triangleright L \triangleright M \triangleright I \triangleright P \triangleright Q \triangleright \{1\}$$

such that the group G/M lies in $\text{Jor}_\ell(J'(n))$, and the group M lies in $\Sigma_\ell(2^n)$ – for the latter use that Q is pro- ℓ and normal in G and P/Q is a finite ℓ -group. $\square$

The following lemma records a useful permanence property of groups in $\Sigma_\ell(c)$ and $\text{Jor}_\ell(d)$.

Lemma 3.9. *Fix any $e \in \mathbb{N}$. Then for any prime number ℓ the following holds:*

- (a) *If $H \trianglelefteq H$ is a normal subgroup of some $H \in \text{Jor}_\ell(e)$, then H' and H/H' lie in $\text{Jor}_\ell(e)$.*
- (b) *If $M' \trianglelefteq M$ is a closed normal subgroup of some $M \in \Sigma_\ell(e)$, then M' and M/M' lie in $\Sigma_\ell(e)$.*

If M' in part (b) of the lemma was not normal in M then clearly M' need not lie in $\Sigma_\ell(c)$ again.

Proof. We only give the proof of (b), the proof of (a) being similar but simpler. Let M be in $\Sigma_\ell(e)$ and consider a normal series $M \triangleright I \triangleright P \triangleright \{1\}$ as in Definition 3.1. Then $L := M/I$ is isomorphic to a product $L_1 \times \cdots \times L_s$ for certain finite simple groups of Lie type L_i in characteristic ℓ . Suppose M' is a closed normal subgroup of M and define $\overline{M'} = M'I/I$. By Goursat's Lemma the groups $\overline{M'}$ and $M'/\overline{M'}$ are products of some of the L_i . From this it is straightforward to see that both M' and M/M' lie in $\Sigma_\ell(c)$. $\square$

The following corollary is immediate from Lemma 3.9(b):

Corollary 3.10. *Fix a constant $c \in \mathbb{N}$. Let G be a profinite group, and for each $\ell \in L$ let $\rho_\ell: G \rightarrow G_\ell$ be a homomorphism of profinite groups such that $\text{Im}(\rho_\ell) \in \Sigma_\ell(c)$ for all $\ell \in L$. Then for any closed normal subgroup $N \trianglelefteq G$ one has $\text{Im}(\rho_{\ell|_H}) \in \Sigma_\ell(c)$ for all $\ell \in L$.*

In particular, if $H \trianglelefteq G$ is an open subgroup, then the above applies to any normal open subgroup $N \trianglelefteq G$ that is contained in H .

4 Fundamental groups: finiteness properties and ramification

The purpose of this section is to recall some finiteness properties of fundamental groups and to provide some basic results on ramification. Regarding the latter we draw from results by Kerz-Schmidt and Wiesend (cf. [24]) and from de Jong on alterations (cf. [6]).

We begin with a finiteness result of which a key part is from [14].

Proposition 4.1. *Suppose that either k is a finite field and S is a smooth proper k -variety or that k is a number field and S is a smooth k -variety, and denote by $K = k(S)$ be the function field of S . For $d \in \mathbb{N}$, let $\mathcal{M}_d$ be the set of all finite Galois extensions E/K inside $\tilde{K}$ such that $\text{Gal}(E/K)$ satisfies $\text{Jor}(d)$ and such that E is unramified along S . Then there exists a finite Galois extension K'/K which is unramified along S such that $E \subset \tilde{k}K'$ for every $E \in \mathcal{M}_d$.*

Proof. Let $\Omega = \prod_{E \in \mathcal{M}_d} E$ be the compositum of all the fields in $\mathcal{M}_d$. For every $E \in \mathcal{M}_d$ the group $\text{Gal}(E/K)$ satisfies $\text{Jor}(d)$ and hence there is a finite Galois extension E'/K inside E such that $[E' : K] \leq d$ and such that E/E' is abelian. Define

$$\Omega' = \prod_{E \in \mathcal{M}_d} E'.$$

Then Ω/Ω' is abelian. Let κ (resp. k_0) be the algebraic closure of k in Ω (resp. in K). It suffices to prove the following

Claim. The extension $\Omega/\kappa K$ is finite.

In fact, once this is shown, it follows that the finite separable extension $\Omega/\kappa K$ has a primitive element ω . Then $\Omega = \kappa K(\omega)$ and $K(\omega)/K$ is a finite separable extension. Let K' be the normal closure of $K(\omega)/K$ in Ω . Then $\tilde{k}K' \supset \kappa K' \supset \kappa K(\omega) = \Omega$ as desired.

In the case $k = \mathbb{Q}$ the claim has been shown in [14, Proposition 2.2]. Assume from now on that k is finite. It remains to prove the claim in that case. The structure morphism $S \rightarrow \text{Spec}(k)$ of the smooth scheme S factors through $\text{Spec}(k_0)$ and S is a geometrically connected k_0 -variety. The profinite group $\pi_1(S \times_{k_0} \text{Spec}(\tilde{k}))$ is topologically finitely generated (cf. [18, Thm. X.2.9]) and $\text{Gal}(k_0) \cong \hat{\mathbb{Z}}$. Thus it follows by the exact sequence (cf. [18, Thm. IX.6.1])

$$1 \rightarrow \pi_1(S \times_{k_0} \text{Spec}(\tilde{k})) \rightarrow \pi_1(S) \rightarrow \text{Gal}(k_0) \rightarrow 1$$

that $\pi_1(S)$ is topologically finitely generated. Thus there are only finitely many extensions of K in $\tilde{K}$ of degree $\leq d$ which are unramified along S . It follows that Ω'/K is a *finite* extension. If we denote by S' the normalization of S in Ω' , then $S' \rightarrow S$ is finite and étale, hence S'/k is a smooth proper variety. Furthermore Ω/Ω' is abelian and unramified along S' . Hence $\Omega/\kappa\Omega'$ is finite by Katz-Lang (cf. [22, Thm. 2, p. 306]). As Ω'/K is finite, it follows that $\Omega/\kappa K$ is finite. $\square$

Our next aim is to introduce several notions of ramification, that are refinements of [24], useful for coverings of general schemes. We fix some terminology for curves. A curve C over a field k is a smooth (but not necessarily projective) k -variety of dimension 1. By $k(C)$ we denote its function field, and by $P(C)$ the smooth projective model of $k(C)$ (which is unique up to isomorphism). Note that $P(C)$ contains C , and we set $\partial C := P(C) \setminus C$. If $\text{char}(k) = p \neq \ell$, then an étale cover $C' \rightarrow C$ is ℓ -*tame* if for any point $x \in \partial C$ with valuation v_x of $k(C)$, the extension $k(C')/k(C)$ is at most tamely ramified² and for any point y of $\partial C'$ above x , the ramification index of y over x is a power of ℓ . The cover $C' \rightarrow C$ is *tame* if the extension $k(C')/k(C)$ is at most tamely ramified.

Definition 4.2. Let $f: T \rightarrow S$ be an étale morphism of regular varieties over a field k of characteristic $p \geq 0$. Let ℓ be a prime different from p and denote by E and K the function fields of T and S , respectively. We define:

²This includes the requirement that all extensions of all residue fields are separable.

- (a) The cover T/S (given by f) is curve ℓ -tame if for all k -morphisms $\varphi: C \rightarrow S$ with C a smooth curve over k , the base changed cover $f_C: C \times_S T \rightarrow C$ is ℓ -tame. (The cover is curve-tame if for all φ the cover f_C is tame.)
- (b) The extension E/K is divisor ℓ -tame if for all discrete rank 1 valuations v of K and w of E above v , the extension E_w/K_v of the completions at w and v , respectively, is tame and the ramification indices are powers of ℓ . (The extension E/K is divisor tame if for all discrete rank 1 valuations v the extension E_w/K_v is tame.)
- (c) If S is an open subscheme of a regular projective scheme $\overline{S}$ such that $D = \overline{S} \setminus S$ is a normal crossings divisor (NCD), then f is ℓ -tame if condition (b) holds for all valuations defined by the generic points of D . (The cover is tame if the extension E/K is divisor tame at all valuations defined by the generic points of D .)

We extend the above three notions to profinite étale covers by saying that such a cover is pro- ℓ curve tame, pro- ℓ divisor tame or pro- ℓ tame if these conditions holds for all subcovers of finite degree.

We note right away that if in the above definition $T \rightarrow S$ is finite étale and Galois, then condition (b) is equivalent to the following condition (cf. [24, Rem. 3.4])

- (b') For all discrete rank 1 valuations v of K and w of E above v , the ramification group I_w inside $\text{Gal}(E_w/K_v)$ is an ℓ -group.³

Note also that since ℓ is different from p , condition (b') implies that the residue extension of E_w/K_v is separable.

Definition 4.3. Suppose k is a field and K/k is a finitely generated extension. We call a homomorphism $\text{Gal}(K) \rightarrow G$ of profinite groups ℓ -tame over k if:

- (a) it factors via $\pi_1(S)$ for some regular variety S over k with function field $k(S) = K$ and
- (b) the fixed field $E := (K_s)^{\text{Ker}(\rho)}$ is pro- ℓ divisor tame over K .

Remark 4.4. In [24, p. 12] $f: T \rightarrow S$ is defined to be *divisor tame* if for any normal compactification $\overline{S}$ of S and a point $s \in \overline{S} \setminus S$ with $\text{codim}_{\overline{S}} s = 1$, the rank 1 valuation v_s on K defined by s is tamely ramified in E/K . We claim that this condition is equivalent to the one we give (assuming that $K \rightarrow E$ is generically étale via some f):

Clearly our notion of divisor tameness implies that of [24]. For the converse we follow closely the argument in [24, Thm. 4.4] proof of (ii) $\Rightarrow$ (iii) though with different references. Let w be a valuation of E that is trivial on k and denote by v its restriction to K . Let $\overline{S}$ be a normal compactification of S , which exists by the theorem of Nagata [28]. By [37, Prop. 6.4], there exists a blow-up $\overline{S}'$ of $\overline{S}$ with center outside S such that v is the valuation of a codimension 1

³If $k(w)$ and $k(v)$ denote the respective residue fields, then I_w is defined as the kernel of the canonical homomorphism $\text{Aut}(E_w/K_v) \rightarrow \text{Aut}(k(w)/k(v))$.

point $s \in \overline{S}'$. By normalization, we may further assume that $\overline{S}'$ is normal. Both operations, blow-up and normalization, do not affect S , and so we may take for $\overline{S}$ a normal compactification of S that contains a codimension 1 point s with valuation $v = v_s$. Because $E = k(\overline{T})$, we have $w = v_t$ for some codimension 1 point of T above s . But then the divisor tameness of [24] implies that w/v is at most tamely ramified.

The following result is a variant of parts of [24, Thm. 4.4]:

Proposition 4.5. *Let k, S, T, K, E, f be as in Definition 4.2. Then the following hold:*

- (a) *The cover T/S is curve-tame if and only if the extension E/K is divisor tame.*
- (b) *Suppose E/K is Galois. Then the cover T/S is curve- ℓ -tame if and only if the extension E/K is divisor ℓ -tame.*
- (c) *If S is an open subscheme of a regular projective scheme $\overline{S}$ such that $D = \overline{S} \setminus S$ is a NCD, then both conditions from (b) are equivalent to T/S being ℓ -tame.*

The assertions (a)–(c) extend in an obvious manner to profinite covers.

Proof. In light of Remark 4.4, part (a) of the proposition follows directly from the equivalence (i) $\Leftrightarrow$ (ii) in [24, Thm. 4.4].

For the proof of (b), suppose first that T/S is curve- ℓ -tame and assume that E/K is not divisor ℓ -tame. By (a) we know that E/K is divisor tame. So let w be a valuation of E at which E/K is not divisor ℓ -tame. Denote by $K_1 \subset K_2 \subset E$ extensions of K such that E/K_1 is totally ramified (and Galois) at w and K_2/K_1 is of prime degree $\ell' \neq \ell$. As in Remark 4.4, there exists a normal compactification $\overline{S}$ of S and a codimension 1 point s of $\overline{S} \setminus S$ that has a preimage t in the normalization $\overline{T}$ of $\overline{S}$ in E/K with $v_t = w$. We define $T_i, \overline{T}_i$, $i = 1, 2$, as the normalizations of S or $\overline{S}$ in K_i/K , respectively. We claim that T/T_1 is curve- ℓ -tame.

To see this, observe first, that as with curve-tameness, it is a simple matter of drawing a suitable commutative diagram to see that curve- ℓ -tameness is stable under base change. In particular the base change $T \times_S T_1 \rightarrow T_1$ is curve- ℓ -tame. However, considering the commutative fiber product diagram

$$\begin{array}{ccc} T & \xleftarrow{\quad s \quad} & T \times_S T_1 \\ \downarrow & \searrow & \downarrow \\ S & \xleftarrow{\quad} & T_1, \end{array}$$

we see that there is a canonical splitting $s: T \rightarrow T \times_S T_1$ over T_1 . Hence T is a connected component of $T \times_S T_1$ and as such the restriction $T \rightarrow T_1$ of the morphism $T \times_S T_1 \rightarrow T_1$ inherits curve- ℓ -tameness.

Having the claim at our disposal, the hypothesis $[K_2 : K_1] = \ell'$ yields that for any curve C_1 mapping to T_1 , the induced cover $C_1 \times_{T_1} T_2 \rightarrow C_1$ is everywhere unramified along ∂C_1 . Now $\overline{T}_1$ is regular in codimension 1, hence the regular locus W_1 contains T_1 as well as the divisor corresponding to $w|_{K_1}$. Let W_2 be its preimage in $\overline{T}_2$. Now by [24, Prop. 4.1], which can be

paraphrased as: *curve-unramifiedness implies unramifiedness over a regular base*, it follows that $W_2 \rightarrow W_1$ is étale. But then K_2/K_1 is étale along w , a contradiction.

For the converse of (b) suppose that E/K is divisor ℓ -tame. We assume that there is a k -morphism $C \rightarrow S$ for C a smooth curve such that $\pi: C \times_S T \rightarrow S$ is not ℓ -tame along ∂C . Since $\text{Gal}(E/K)$ acts faithfully on $C \times_S T \rightarrow C$, by passing to a subgroup and thus an intermediate extension of E/K we may assume that $C \times_S T$ is irreducible. Since then $\text{Gal}(E/K)$ is also the Galois group of the cover π , some further straightforward reductions allow us to assume that $[E : K] = \ell' \neq \ell$ for some prime ℓ' (which by (a) is different from p), and that $\text{Gal}(E/K) \cong \mathbb{Z}/\ell'$ is the inertia group above some valuation of $k(C)$. Following the argument in the proof of [24, Thm. 4.4] (v) $\Rightarrow$ (i), we can find a discrete rank $d = \dim S$ valuation of E that is ramified of order ℓ' (via a Parshin chain through the image of $\text{Spec } k(C)$). But [24, Lem. 3.5] says that E/K is ramified at a discrete rank d valuation if and only if it is ramified at a discrete rank 1 valuation. We reach a contradiction because by hypothesis E/K is unramified at all discrete rank 1 valuations.

Finally, we prove (c). It is clear that divisor ℓ -tameness implies ℓ -tameness. The proof that ℓ -tameness implies curve ℓ -tameness follows from that argument given in [24, Prop. 4.2]: there it is shown that tameness implies curve tameness. Consider a curve C over k and a morphism $\varphi: C \rightarrow S$ over k . Then φ extends to a morphism $\bar{\varphi}: P(C) \rightarrow \bar{S}$. Denote by $\bar{\varphi}(C)$ the closure of $\varphi(C)$ in $\bar{S}$. The ramification of $T \times_S C \rightarrow C$ occurs precisely at those points of $P(C)$ that under $\bar{\varphi}$ map to $D \cap \bar{\varphi}(C)$. To analyze the ramification, the proof of [24, Prop. 4.2] appeals to Abhyankar's lemma. In the notation of *loc. cit.*, the ramification is then governed by indices n_i , $i = 1, \dots, r$, that are prime to p . By the ℓ -tameness of $T \rightarrow S$, the n_i must all be powers of ℓ . But then *loc. cit.* implies that $T \times_S C \rightarrow C$ is ℓ -tame, and this completes the proof. $\square$

Our formulation of divisor-tameness easily transfers under rather general field extensions:

Lemma 4.6. *Suppose that $\text{char}(k) = p > 0$ and consider the following inclusions of fields:*

$$\begin{array}{ccc} K & \hookrightarrow & K' \\ \uparrow & & \uparrow \\ k & \hookrightarrow & k' \end{array}$$

If E/K is Galois and divisor ℓ -tame over k (or divisor pro- ℓ tame), then so is EK'/K' over k' .

Proof. It clearly suffices to prove the lemma in the case where E/K is finite Galois. Then $E' := EK'$ is finite Galois over K' . Let w' be any discrete rank one valuation of E' trivial on k' and denote by w its restriction to E , by v' the restriction to K' and by v the restriction to K . We need to show that $[w'(E'^*) : v'(K'^*)]$ is a power of ℓ and that the residue extension is separable. The latter can be taken care of at once: The extension E/K is finite separable. Hence so is E'/K' , because a primitive element of E/K will be such an element for E'/K' . For the same reason, separability is preserved by the extensions of completions $E'_{w'}/K'_{v'}$ at v' . Now by the Cohen structure theorem, the extension of residue fields is a subextension of $E'_{w'}/K'_{v'}$, and as such it must be separable. It remains to consider the index of the value groups.

Suppose first that $v(K^*) = 0$. Then we must have $w(E^*) = 0$, since otherwise, if $\alpha \in E$ would satisfy $w(\alpha) \neq 0$, then the sequence $(\alpha^n)_{n \in \mathbb{Z}}$ would be linearly independent over K , a contradiction. This means that under the residue map of E' , the subfield E is mapped injectively to the residue field of E' at w' . But then E/K defines purely a residue extension of E'/K' , and thus $w'(E^*) = v'(K'^*)$.

Suppose now that w is non-trivial, so that by the above v is non-trivial as well. We pass to the completions and note that E_w/K_v and $E'_{w'}/K_{v'}$ remain Galois extensions. By the Cohen structure theorem, K_v now contains the residue field $k(v)$, and E_v the residue field $k(w)$. In particular, $F = k(w)K_v$ is an unramified extension of K_v and E_w/F is totally ramified. We may thus consider these two cases separately. Suppose first that E_w/K_v is unramified. Then $E'_{w'} = K'_{v'}k(w)$ where clearly $k(w)$ defines a separable extension of the residue field of $K'_{v'}$. Hence $E'_{w'}/K'_{v'}$ is unramified. We conclude $w'(E^*) = v'(K'^*)$ which completes the argument.

Suppose now that E_w/K_v is totally ramified. By our hypothesis, the extension E/K is at most ℓ -order ramified at w . It follows that E_w/K_v is a Galois extension with $\text{Gal}(E_w/K_v)$ an ℓ -group. Now $\text{Gal}(E'_{w'}/K'_{v'})$ injects into $\text{Gal}(E_w/K_v)$ because $E' = KE$, and thus $[E'_{w'} : K'_{v'}]$ is a power of ℓ . Since the order of $w'(E^*)/v'(K'^*)$ divides the degree $[E' : K']$, the proof is complete. $\square$

Combining ramification properties with finiteness properties of fundamental groups, we obtain the following criterion for a family of representations of $\text{Gal}(K)$ with images in $\text{Jor}_\ell(d)$, or with abelian images of bounded order, to become trivial over $\text{Gal}(K'\tilde{k})$ for some finite K'/K .

Proposition 4.7. *Let k be a field and let S/k be a normal k -variety with function field K . Suppose $(\rho_\ell : \pi_1(S) \rightarrow G_\ell)_{\ell \in L}$ is a family of continuous homomorphisms such that if $\text{char}(k) > 0$ each ρ_ℓ is ℓ -tame. Under either of the following two conditions there exists a finite extension K' of K such that for all $\ell \in L$ we have $\rho_\ell(\text{Gal}(K'\tilde{k})) = \{1\}$.*

- (a) *The field k is finite or a number field and there exists a constant $d \in \mathbb{N}$ such that for each $\ell \in L$ the group $\text{Im}(\rho_\ell)$ lies in $\text{Jor}_\ell(d)$.*
- (b) *The field k is algebraically closed and there exists a constant $c \in \mathbb{N}$ such that for each $\ell \in L$ the group $\text{Im}(\rho_\ell)$ is abelian of order at most c .*

Proof. First we replace K by a finite Galois extension and S be the normalization in this extension, so that we can assume that $\rho_\ell(\pi_1(S)) = \{1\}$ for all $\ell \leq d$ or $\ell \leq c$, respectively. Next we apply the result of de Jong on alterations (cf. [6, Thm. 4.1, 4.2]). It provides us with a finite extension k' of k , a smooth projective geometrically connected k' -variety T' , a non-empty open subvariety S' of T' and an alteration $f : S' \rightarrow S$, such that furthermore $D' := T' \setminus S'$ is a normal crossing divisor. We define K' to be the function field of S' , so that K'/K is finite. (If k is perfect, we could also assume that K'/K is separable.)

Next observe, that if $\text{char}(k)$ is positive, then the (divisor) ℓ -tameness of ρ_ℓ implies the (divisor) ℓ -tameness of $\rho_\ell|_{\pi_1(S')}$ by Lemma 4.6, and thus, by Proposition 4.5, for each ℓ the extension $(K'_s)^{\text{Ker}(\rho_\ell|_{\pi_1(S')})}$ of K' is ℓ -tame. Because of the first reduction step in the previous paragraph, this implies that each $\rho_\ell|_{\pi_1(S')}$ is unramified at the generic points of D' . Purity of the branch locus (cf. [18, X.3.1]) now implies that all $\rho_\ell|_{\pi_1(S')}$ factor via $\pi_1(T')$.

Now, in case (a), the assertion of the proposition follows from Proposition 4.1. In case (b) we use that, since k is algebraically closed, the fundamental group $\pi_1(T')$ is topologically finitely generated (cf. [18, Thm. X.2.9]), and that furthermore, if $\text{char}(k) = 0$, the same holds true for $\pi_1(S')$ (cf. [19, II.2.3.1]). Hence in the former case $\pi_1(T')^{\text{ab}}/c\pi_1(T')^{\text{ab}}$ and in the latter case $\pi_1(S')^{\text{ab}}/c\pi_1(S')$ are finite. This completes the proof in case (b). $\square$

5 An independence criterion

From now on, let k be any field, let K/k be a finitely generated field extension and let L be a set of prime numbers not containing $p := \text{char}(k)$. For every $\ell \in L$ let G_ℓ be a profinite group and let $\rho_\ell : \text{Gal}(K) \rightarrow G_\ell$ be a continuous homomorphism.

If for all $\ell \in L$ the groups $\text{Im}(\rho_\ell)$ are n -bounded at ℓ , then by Theorem 3.4 we have a short exact sequence $1 \rightarrow M_\ell \rightarrow \text{Im}(\rho_\ell) \rightarrow H_\ell \rightarrow 1$ with $H_\ell \in \text{Jor}_\ell(d)$ for $d = J'(n)$ and $M_\ell \in \Sigma_\ell(2^n)$. At the end of the previous section we have seen that a combination of tameness of ramification and results on fundamental groups allow one to control the H_ℓ in a uniform manner. In this section we shall show how to control M_ℓ in a uniform manner, if one has a uniform control on ramification. We begin by introducing the necessary concepts and then give the result.

To $(\rho_\ell)_{\ell \in L}$ we attach the family $(\tilde{\rho}_\ell)_{\ell \in L}$ by defining each $\tilde{\rho}_\ell$ as the composite homomorphism

$$\tilde{\rho}_\ell : \text{Gal}(K) \xrightarrow{\rho_\ell} \text{Im}(\rho_\ell) \rightarrow \text{Im}(\rho_\ell)/Q_\ell$$

where Q_ℓ is the maximal normal pro- ℓ subgroup of $\text{im}(\rho_\ell)$. Note that if ρ_ℓ is an ℓ -adic representation, then $\tilde{\rho}_\ell$ is essentially the semisimplification of the mod ℓ reduction of ρ_ℓ .

Definition 5.1. (a) The family $(\rho_\ell)_{\ell \in L}$ is said to satisfy the condition $\mathcal{R}(k)$, if there exist a finite extension k' of k , a finite extension K'/Kk' and a smooth k' -variety U' with function field K' such that for every $\ell \in L$ the homomorphism $\tilde{\rho}_\ell|_{\text{Gal}(K')}$ is unramified along U' .

(b) The family $(\rho_\ell)_{\ell \in L}$ is said to satisfy the condition $\mathcal{S}(k)$, if it satisfies $\mathcal{R}(k)$ and if one can choose the field K' for $\mathcal{R}(k)$ such that each $\tilde{\rho}_\ell|_{\text{Gal}(K')}$ is ℓ -tame.

The condition $\mathcal{R}(k)$ says that each member $\tilde{\rho}_\ell$ is up to pro- ℓ ramification potentially generically étale in a uniform way. The condition $\mathcal{S}(k)$ is a kind of semistability condition.

Example 5.2. Set $L = \mathbb{L} \setminus \{\text{char}(k)\}$ and let A/K be an abelian variety. For every $\ell \in L$ denote by $\sigma_\ell : \text{Gal}(K) \rightarrow \text{Aut}_{\mathbb{Z}_\ell}(T_\ell(A))$ the representation of $\text{Gal}(K)$ on the ℓ -adic Tate module $\varprojlim_{i \in \mathbb{N}} A[\ell^i]$. There exists a finite extension k' of k and a finite separable extension $K'/k'K$ such that K' is the function field of some smooth k' -variety V' . By the spreading-out principles of [17] there exists a non-empty open subscheme U' of V' and an abelian scheme $\mathcal{A}$ over U' with generic fibre A . This implies (cf. [19, IX.2.2.9]) that σ_ℓ is unramified along U' for every $\ell \in L$. Hence the family $(\sigma_\ell)_{\ell \in L}$ satisfies condition $\mathcal{R}(k)$.

In order to obtain also $\mathcal{S}(k)$, we choose an odd prime $\ell_0 \in L$, and we require the field K' above to be finite separable over $k'K(A[\ell_0])$. Now let v' be any discrete valuation of K' which is trivial

on k' , and let $R_{v'}$ be the discrete valuation ring of v' . Let $\mathcal{N}_{v'}/\mathrm{Spec}(R_{v'})$ be the Néron model of A over $R_{v'}$. The condition $K' \supset K(A[\ell_0])$ forces $\mathcal{N}_{v'}$ to be semistable (cf. [19, IX.4.7]). This in turn implies that σ_ℓ is unipotent (and hence pro- ℓ) for every $\ell \in L$ (cf. [19, IX.3.5]). It follows that the family $(\sigma_\ell)_{\ell \in L}$ satisfies condition $\mathcal{S}(k)$.

The following is the main independence criterion of this section:

Proposition 5.3. *Let k be an algebraically closed field and let K/k be a finitely generated extension. Fix a constant $c \in \mathbb{N}$ and suppose that $(\rho_\ell)_{\ell \in L}$ is a family of representations of $\mathrm{Gal}(K)$ that satisfies the following conditions:*

- (a) *The family $(\rho_\ell)_{\ell \in L}$ satisfies $\mathcal{R}(k)$ if $\mathrm{char}(k) = 0$ and $\mathcal{S}(k)$ if $\mathrm{char}(k) > 0$.*
- (b) *There exists a constant $c \in \mathbb{N}$ such that for all $\ell \in L$ one has $\mathrm{Im}(\rho_\ell) \in \Sigma_\ell(c)$.*

Then there exists a finite abelian Galois extension E/K with the following properties.

- (i) *For every $\ell \in L$ the group $\rho_\ell(\mathrm{Gal}(E))$ lies in $\Sigma_\ell(c)$ and is generated by its ℓ -Sylow subgroups, and for $\ell > c$ it is generated by the ℓ -Sylow subgroups of $\rho_\ell(\mathrm{Gal}(K))$.*
- (ii) *The group $\mathrm{Gal}(E)$ is a characteristic subgroup of $\mathrm{Gal}(K)$.*
- (iii) *The restricted family $(\rho_\ell|_{\mathrm{Gal}(E)})_{\ell \in L \setminus \{2,3\}}$ is independent and $(\rho_\ell)_{\ell \in L}$ is almost independent.*

Proof. We can assume that ρ_ℓ is surjective for all $\ell \in L$. Denote by G_ℓ^+ the normal subgroup of G_ℓ which is generated by the pro- ℓ Sylow subgroups of G_ℓ . Then $\overline{G}_\ell := G_\ell/G_\ell^+$ is a finite group of order prime to ℓ . As G_ℓ lies in $\Sigma_\ell(c)$, so does its quotient $\overline{G}_\ell$ by Lemma 3.9(b). Now any group in $\Sigma_\ell(c)$ of order prime to ℓ is abelian of order at most c , and thus the latter holds for $\overline{G}_\ell$.

Let K_ℓ^+ be the fixed field in K_s of the kernel of the map $\pi_\ell \circ \rho_\ell$, so that $\mathrm{Gal}(K_\ell^+/K) \cong \overline{G}_\ell$. Then the compositum $E = \prod_{\ell \in L} K_\ell^+$ is an abelian extension of K such that $\mathrm{Gal}(E/K)$ is annihilated by $c!$. From Proposition 4.7(b) we see that E/K is finite. Assertion (ii) is now straightforward: By definition of the K_ℓ^+ the subgroups $\mathrm{Gal}(K_\ell^+)$ of $\mathrm{Gal}(K)$ are characteristic and hence so is their intersection $\mathrm{Gal}(E)$.

We turn to the proof of (i): For every $\ell \in L$, from (ii) the group $\rho_\ell(\mathrm{Gal}(E))$ is normal in G_ℓ , and hence it lies in $\Sigma_\ell(c)$ by Lemma 3.9. By construction $\rho_\ell(\mathrm{Gal}(E)) \subset \rho_\ell(\mathrm{Gal}(K_\ell^+)) = G_\ell^+$ and $N_\ell := G_\ell^+/\rho_\ell(\mathrm{Gal}(E))$ is abelian and annihilated by $c!$. We claim that (1) N_ℓ is an ℓ -group, so that N_ℓ is trivial if $\ell > c$, and that (2) $\rho_\ell(\mathrm{Gal}(E))$ is generated by its pro- ℓ Sylow subgroups. We argue by contradiction and assume that (1) or (2) fails.

If (2) fails, then $\rho_\ell(\mathrm{Gal}(E))$ has a finite simple quotient of order prime to ℓ . Because $\rho_\ell(\mathrm{Gal}(E))$ lies in $\Sigma_\ell(c)$, this simple quotient has to be abelian of prime order ℓ' different from ℓ . Again by (ii), the Galois closure over K of the fixed field of this ℓ' -extension is a solvable extension. Denote by F either this solvable extension if (2) fails, or the extension of K_ℓ^+ whose Galois group is canonically isomorphic to N_ℓ if (1) fails. In either case F/K is Galois and solvable, and we have a canonical surjection $\pi'_\ell: G_\ell \twoheadrightarrow \mathrm{Gal}(F/K)$. Arguing as in the first paragraph,

it follows that I_ℓ surjects onto $\text{Gal}(F/K)$. By construction $\text{Gal}(F/K_\ell^+)$ is not an ℓ -group. It follows from the definition of K_ℓ^+ that the normal subgroup $\pi'_\ell(P_\ell) \subset \text{Gal}(F/K)$ is a proper subgroup of $\text{Gal}(F/K_\ell^+)$. But then its fixed field is a proper extension of K_ℓ^+ which is at once Galois and of a degree over K that is prime to ℓ . This contradicts the definition of K_ℓ^+ , and thus (1) and (2) hold. This in turn completes the proof of (i).

We now prove (iii). Denote by Ξ_ℓ the class of those finite groups which are either a finite simple group of Lie type in characteristic ℓ or isomorphic to $\mathbb{Z}/(\ell)$. The conditions in (i) imply that every simple quotient of $\rho_\ell(\text{Gal}(E))$ lies in Ξ_ℓ . But now for any $\ell, \ell' \geq 5$ such that $\ell \neq \ell'$ one has $\Xi_\ell \cap \Xi_{\ell'} = \emptyset$ (cf. [35, Thm. 5], [1], [25]). The first part of (iii) now follows from [35, Lemme 2]. The second part follows from the first, the definition of almost independence and from [35, Lemme 3]. $\square$

6 Reduction steps

Let k, K, L, p and $(\rho_\ell)_{\ell \in L}$ be as at the beginning of Section 5. In the previous two sections we have described ramification properties of $(\rho_\ell)_{\ell \in L}$ and properties of $(\rho_\ell(\text{Gal}(K)))_{\ell \in L}$ that were essential to control in a uniform way the groups H_ℓ and M_ℓ that occur in $\rho_\ell(\text{Gal}(K))$ as in Theorem 3.4. The aim of this section is to explain how these properties for a general pair (K, k) in our target Theorem 1.2, can be reduced to a pair where k is the prime field and K is finitely generated over it. Moreover we shall explain how one can reduce the proof of our target theorem to the case where X is a smooth and projective variety over K .

Lemma 6.1. *Suppose we have a commutative diagram of fields*

$$\begin{array}{ccc} K & \hookrightarrow & K' \\ \uparrow & & \uparrow \\ k & \hookrightarrow & k' \end{array}$$

such that K' is finite over Kk' . Then the following properties hold true:

- (i) *If $(\rho_\ell)_{\ell \in L}$ satisfies $\mathcal{R}(k)$, then $(\rho_\ell|_{\text{Gal}(K')})_{\ell \in L}$ satisfies $\mathcal{R}(k')$.*
- (ii) *If $\text{char}(k) > 0$ and $(\rho_\ell)_{\ell \in L}$ satisfies $\mathcal{S}(k)$, then $(\rho_\ell|_{\text{Gal}(K')})_{\ell \in L}$ satisfies $\mathcal{S}(k')$.*
- (iii) *If there exists a constant $c \in \mathbb{N}$ such that for all $\ell \in L$, the group $\rho_\ell(\text{Gal}(K\tilde{k}))$ lies in $\Sigma_\ell(c)$, then there exists a finite Galois extension E'/K' such that for all $\ell \in L$, the group $\rho_\ell(\text{Gal}(E'\tilde{k}'))$ lies in $\Sigma_\ell(c)$.*


Proof. Considering the diagram

$$\begin{array}{ccccccc} K & \xlongequal{\quad} & K & \hookrightarrow & k'K & \hookrightarrow & K' \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ k & \hookrightarrow & k' \cap K & \hookrightarrow & k' & \xlongequal{\quad} & k', \end{array}$$

it suffices to prove the lemma in the following three particular cases: (a) $K' = K$, (b) $Kk' = K'$ and $K \cap k' = k$ (base change), (c) $k = k'$ and K' is finite over K .

For the proof of (iii) note that in case (a) the group $\text{Gal}(K'\tilde{k}')$ is a closed normal subgroup of $\text{Gal}(K\tilde{k})$ and in case (b) the canonical homomorphism $\text{Gal}(K'\tilde{k}') \rightarrow \text{Gal}(K\tilde{k})$ is surjective. In either case we take $E = K'$. Finally in case (c) we define E as the Galois closure of K' over K . Then in cases (a) and (c) assertion (iii) follows from Corollary 3.10 with $H = \text{Gal}(E)$ and $G = \text{Gal}(K)$. Case (b) is obvious.

In all cases, the proof of (ii) is immediate from Lemma 4.6, once (i) is proved. Therefore it remains to prove (i). We first consider case (a). By replacing k , k' and K several times by finite extensions, we can successively achieve the following, where in each step the previous property is preserved: First, using de Jong's result on alterations (cf. [6, Thm. 4.1, 4.2]), there exists a smooth projective scheme X/k' whose function field is K . Second, by the spreading out principle, there exists an affine scheme U' over k whose function field is k' and a smooth projective U' -scheme $\mathcal{X}$ whose function field is K . Third, by hypothesis $\mathcal{R}(k)$ there exists a smooth k -scheme U whose function field is K such that all $\tilde{\rho}_\ell$ factor via $\pi_1(U)$. By shrinking U we may assume it to be affine. Also we choose an affine open subscheme $\mathcal{V}$ of $\mathcal{X}$. The corresponding coordinate rings we denote by R and $\mathcal{R}$, respectively. Both of these rings are finitely generated over k . Since the fraction field of both is K , by inverting a suitable element $g \neq 0$ of R we have $R[g^{-1}] \supset \mathcal{R}$, and similarly we can find $0 \neq f \in \mathcal{R}$ such that $\mathcal{R}[f^{-1}] \supset R$. Inverting both elements shows that we can find an affine open subscheme V of both U and $\mathcal{V}$. In particular, the function field of V is K , the scheme V is smooth over k and over U' and the representations ρ_ℓ all factor via $\pi_1(V)$. The following diagram displays the situation:



Define V' as the base change $V \times_{U'} \text{Spec } k'$, so that V' to $\text{Spec } k'$ is smooth and affine. Now if $W \rightarrow U$ is any étale Galois cover, then the base change $W \times_U V'$ is an étale Galois cover of V' . We deduce that $\tilde{\rho}_\ell$ factors via $\pi_1(V')$ for all ℓ , and thus we have verified $\mathcal{R}(k')$ for $(\tilde{\rho}_\ell)_{\ell \in L}$.

Case (b): This is a base change. Therefore if we replace k by a finite extension, then U/k becomes a smooth variety with function field a finite separable extension of Kk' . But then $U \times_{\text{Spec } k} \text{Spec } k'$ is smooth over k' and its function field is a finite separable extension of $K' = Kk'$. From this (i) is immediate.

Case (c): To see $\mathcal{R}(k)$ over K' , let $k' \supset k$ and $K'' \supset K'k'$ be finite extensions such that there exists a smooth k' -scheme U with function field K'' such that all $\tilde{\rho}_\ell$ factor via $\pi_1(U)$. Let U' be

the normalization of U in $K'K''$. Now choose $k'' \supset k'$ and $K''' \supset K'K''$ finite such that there is a smooth k'' -scheme U'' with function field K'' and a finite morphism to U' . Then $\mathcal{R}(k')$ is verified by U'' . $\square$

The following is a standard lemma from algebraic geometry about models of schemes over finitely generated fields.

Lemma 6.2. *Let k be a field, K/k be finitely generated and X be an separated algebraic scheme over K . Then there exists an absolutely finitely generated field $K_0 \subset K$ and a separated algebraic scheme X_0 over K_0 such that $kK_0 = K$ and $X_0 \otimes_{K_0} K = X$. If in addition X/K is smooth and projective, then one can choose X_0 and K_0 in a way so that X_0/K_0 is smooth projective.*

Proof. Let $\mathfrak{K}$ be the set of all finitely generated subfields of K . Then $K = \bigcup_{K' \in \mathfrak{K}} K'$ and $\text{Spec}(K) = \varprojlim_{K' \in \mathfrak{K}} \text{Spec}(K')$. There exists $K' \in \mathfrak{K}$ and a separated algebraic K' -scheme X_0 such that $X = X_{0,K'}$ (cf. [17, 8.8.2] and [17, 8.10.5(v)]). If X/K is projective, then one can choose K' and X_0 in such a way that X_0/K' is projective (cf. [17, 8.10.5(xiii)]). If X/K is smooth, then X_0/K' is smooth. Furthermore there exist $x_1, \dots, x_t \in K$ such that $K = k(x_1, \dots, x_t)$. Define $K_0 := K'(x_1, \dots, x_t)$. Then $kK_0 = K$, the field K_0 is finitely generated and $X_0 := X_{0,K_0}$ has the desired properties. $\square$

For a separated algebraic scheme X over K and any $\ell \in \mathbb{L} \setminus \{\text{char}(k)\}$ we denote by $\rho_{\ell,X}$ the representation of $\text{Gal}(K)$ on $\bigoplus_{q \geq 0} (H_c^q(X_{\tilde{K}}, \mathbb{Q}_\ell) \oplus H^q(X_{\tilde{K}}, \mathbb{Q}_\ell))$.

Corollary 6.3. *Suppose that for all absolutely finitely generated fields K_0 with field of constants k_0 and for all schemes X_0 that are separated algebraic over K_0 , the following conditions are true:*

- (a) *The family $(\rho_{\ell,X_0})_{\ell \in L}$ satisfies $\mathcal{R}(k_0)$ if $\text{char}(k_0) = 0$ and $\mathcal{S}(k_0)$ if $\text{char}(k_0) > 0$.*
- (b) *There exists a constant $c \in \mathbb{N}$ and a finite extension E_0 of K_0 such that for all $\ell \in L$ one has $\rho_{\ell,X_0}(\text{Gal}(E_0 \tilde{k}_0)) \in \Sigma_\ell(c)$.*

Let k, K, X be as in Theorem 1.2. Then there exists a finite extension E/K such that the assertions and conclusions of Proposition 5.3 hold for $E\tilde{k}$, and in particular Theorem 1.2 holds.

Proof. Let k be any field, let K over k be a finitely generated extension field and let X be a separated algebraic scheme over K . By Lemma 6.2, we can find $K_0 \subset K$ absolutely finitely generated and X_0 a separated algebraic scheme over K_0 such that $X = X_0 \otimes_{K_0} K$, and moreover if X is smooth and/or projective over K , then the same can be assumed for X_0 over K_0 . Next we replace K_0 by E_0 as guaranteed by our hypotheses and K by $E := E_0 K$. Now Lemma 6.1 yields a finite Galois extension E'/E such that $(\rho_{\ell,X}|_{\text{Gal}(E'\tilde{k})})_{\ell \in L}$ satisfies $\mathcal{R}(\tilde{k})$ if $\text{char}(k) = 0$ and $\mathcal{S}(\tilde{k})$ if $\text{char}(k) > 0$ and in addition that for all $\ell \in L$, the image $\rho_\ell(\text{Gal}(E'\tilde{k}))$ lies in $\Sigma_\ell(c)$. By Proposition 5.3, there exists a finite extension E'' of E' such that the assertions and conclusions of Proposition 5.3 hold for $E''\tilde{k}$. $\square$

We now come to the second reduction step.

Definition 6.4. For a representation $\rho_\ell: \text{Gal}(K) \rightarrow \text{GL}_n(\mathbb{Q}_\ell)$ we denote by ρ_ℓ^{sss} its strict semisimplification, i.e., the direct sum over the irreducible subquotients of ρ_ℓ where each isomorphism type occurs with multiplicity one.

Note that $\text{Im}(\rho_\ell^{\text{sss}}) = \text{Im}(\rho_\ell^{\text{ss}})$ where ρ_ℓ^{ss} denotes the usual semisimplification of ρ_ℓ .

Lemma 6.5. For every $\ell \in L$ let ρ_ℓ and ρ'_ℓ be representations $\text{Gal}(K) \rightarrow \text{GL}_n(\mathbb{Q}_\ell)$. Suppose that one of the following two assertions is true:

- (a) $\rho_\ell^{\text{sss}} = (\rho'_\ell)^{\text{sss}}$ for all $\ell \in L$, or
- (b) ρ_ℓ is a direct summand of ρ'_ℓ for all $\ell \in L$.

Then the following hold:

- (i) if the family $(\rho'_\ell)_{\ell \in L}$ satisfies $\mathcal{R}(k)$ then so does $(\rho_\ell)_L$;
- (ii) if the family $(\rho'_\ell)_{\ell \in L}$ satisfies $\mathcal{S}(k)$ then so does $(\rho_\ell)_L$;
- (iii) for any $\ell \in L$, if $\rho'_\ell(\text{Gal}(K\tilde{k}))$ lies in $\Sigma_\ell(c)$ then so does $\rho_\ell(\text{Gal}(K\tilde{k}))$.

Note that condition (a) is symmetric, so that under (a) each of (i)–(iii) is an equivalence.

Proof. The proof of (i)–(iii) under hypothesis (a) is an immediate consequence of the simple fact that the kernel of $\text{Im}(\rho_\ell) \rightarrow \text{Im}(\rho_\ell^{\text{ss}}) = \text{Im}(\rho_\ell^{\text{sss}})$ is a pro- ℓ -group. Assertions (i) and (ii) under hypothesis (b) are trivial. For (iii) note that hypothesis (b) implies that $\rho'_\ell(\text{Gal}(K\tilde{k}))$ is a closed normal subgroup of $\rho_\ell(\text{Gal}(K\tilde{k}))$, and so we can apply Lemma 3.9. $\square$

The following important result is taken from Seminaire Bourbaki talk of Berthelot on de Jong's alteration technique (cf. [4, Thm. 6.3.2])

Theorem 6.6. Let k be a field and K be a finitely generated extension field. Let X be a separated algebraic scheme over K . Then there exists a finite extension k'/k , a finite separable extension K'/Kk' and a finite set of smooth projective varieties $\{Y_i\}_{i=1,\dots,k}$ over K' such that for all $\ell \in \mathbb{L} \setminus \{\text{char}(k)\}$ the representation $(\rho_{\ell,X}|_{\text{Gal}(K')})^{\text{sss}}$ is a direct summand of $(\bigoplus_i \rho_{\ell,Y_i})^{\text{sss}}$.

Before giving the proof of Theorem 6.6, let us state the following immediate consequence of Theorem 6.6, Lemma 6.5 and Corollary 6.3:

Corollary 6.7. Suppose that for all absolutely finitely generated fields K_0 with field of constants k_0 and for all smooth projective K_0 -varieties X_0/K_0 , the following conditions are true:

- (a) The family $(\rho_{\ell,X_0})_{\ell \in L}$ satisfies $\mathcal{R}(k_0)$ if $\text{char}(k_0) = 0$ and $\mathcal{S}(k_0)$ if $\text{char}(k_0) > 0$.
- (b) There exists a constant $c \in \mathbb{N}$ and a finite extension E_0 of K_0 such that for all $\ell \in L$ one has $\rho_{\ell,X_0}(\text{Gal}(E_0\tilde{k}_0)) \in \Sigma_\ell(c)$.

Let k, K, X be as in Theorem 1.2. Then there exists a finite extension E/K such that the assertions and conclusions of Proposition 5.3 hold for $E\tilde{K}$, and in particular Theorem 1.2 holds.

Proof of Theorem 6.6. For completeness we provide details of the proof in [4]. For $? \in \{c, \emptyset\}$ we denote by $\rho_{\ell, X, ?}$ the representation of $\text{Gal}(K)$ on $\bigoplus_{q \geq 0} (H^q_?(X_{\tilde{K}}, \mathbb{Q}_\ell))$. It suffices to prove the theorem separately for the families $(\rho_{\ell, X, ?})_{\ell \in L}$. We also note that whenever it is convenient, we are allowed (by passing from K to a finite extension) to assume that X is geometrically reduced over K . This is so because $H^q_?(X_{\tilde{K}}, \mathbb{Q}_\ell) \cong H^q_?(X_{\tilde{K}, \text{red}}, \mathbb{Q}_\ell)$ for any $q \in \mathbb{Z}$ and $? \in \{\emptyset, c\}$. We first consider the case of cohomology with compact supports. The proof proceeds by induction on $\dim X$.

We may assume, by passing from K to a finite extension, that X is geometrically reduced. Then $X_{\tilde{K}}$ is generically smooth. After passing from K to a finite extension, we can find a dense open subscheme $U \subset X$ that is smooth over K . By the long exact cohomology sequence with supports (cf. [29, Rem. III.1.30]) we have for any ℓ an exact sequence

$$\dots \longrightarrow H_c^i(U_{\tilde{K}}, \mathbb{Q}_\ell) \longrightarrow H_c^i(X_{\tilde{K}}, \mathbb{Q}_\ell) \longrightarrow H_c^i((X \setminus U)_{\tilde{K}}, \mathbb{Q}_\ell) \longrightarrow \dots,$$

so that for all ℓ the representation $(\rho_{\ell, X, c})_{\ell \in L}^{\text{sss}}$ is a direct summand of $(\rho_{\ell, U, c})_{\ell \in L}^{\text{sss}} \oplus (\rho_{\ell, X \setminus U, c})_{\ell \in L}^{\text{sss}}$. By induction hypothesis, it thus suffices to treat the case that U is smooth over K . By the induction hypothesis it is also sufficient to replace U by any smaller dense open subscheme, and it is clearly also sufficient to treat the case where U is in addition geometrically irreducible.

By de Jong's theorem on alterations (cf. [6, Thms. 4.1, 4.2]), after passing from K to a finite extension, we can find a smooth projective scheme Y , an open subscheme U' of Y and an alteration $\pi: U' \rightarrow U$. By replacing K yet another time by a finite extension, we can assume that $U' \rightarrow U$ is generically finite étale. And now we pass to an open subscheme V of U and to $V' := \pi^{-1}(V) \subset U'$ such that $V' \rightarrow V$ is finite étale. By the induction hypothesis applied to $Y \setminus V'$ and again the long exact cohomology sequence for cohomology with support, we find that the assertion of the theorem holds true for $(\rho_{\ell, V', c})_{\ell \in L}^{\text{sss}}$. From now on π denotes the restriction to V' and $\mathbb{Q}_{\ell, X}$ will be the constant sheaf $\mathbb{Q}_\ell$ on any scheme X . Since π is finite étale, say of degree d , there exists a trace morphism $\text{Trace}_\pi: \pi_* \mathbb{Q}_{\ell, V'} \rightarrow \mathbb{Q}_{\ell, V}$ whose composition with the canonical morphism $\mathbb{Q}_{\ell, V} \rightarrow \pi_* \mathbb{Q}_{\ell, V'}$ is multiplication by d (cf. [29, Lem. V.1.12]). In particular, the constant sheaf $\mathbb{Q}_{\ell, V}$ is a direct summand of $\pi_* \mathbb{Q}_{\ell, V'}$. Since $H_c^i(V'_{\tilde{K}}, \mathbb{Q}_\ell) \cong H_c^i(V_{\tilde{K}}, \pi_* \mathbb{Q}_\ell)$, we deduce that $(\rho_{\ell, V, c})_{\ell \in L}^{\text{sss}}$ is a direct summand of $(\rho_{\ell, V', c})_{\ell \in L}^{\text{sss}}$, and this completes the induction step.

Now we turn to the case $? = \emptyset$. The case when X is smooth over K but not necessarily projective is reduced, by Poincaré duality, to the case of compact supports: one has $H^q(X_{\tilde{K}}, \mathbb{Q}_\ell) \cong H_c^{2d-q}(X_{\tilde{K}}, \mathbb{Q}_\ell(d))^\vee$ for $d = \dim X$ (cf. [29, Cor. VI.11.12]).

Suppose now that X is an arbitrary separated algebraic scheme over K . By what we said above, we may assume that X is geometrically reduced. Again we perform an induction on $\dim X$. The first step is a reduction to the case where X is irreducible, which maybe thought as an induction by itself. Suppose $X = X_1 \cup X_2$ where X_1 is an irreducible component of X and X_2 is the closure of $X \setminus X_1$. Consider the canonical morphism $f: X_1 \sqcup X_2 \rightarrow X$. It yields a short exact sequence of sheaves

$$0 \longrightarrow \mathbb{Q}_{\ell, X} \longrightarrow f_* \mathbb{Q}_{\ell, X_1 \sqcup X_2} \longrightarrow \mathcal{F} \longrightarrow 1 \quad (1)$$

where $\mathcal{F}$ is a sheaf on X . We claim that $\mathcal{F} \cong \mathbb{Q}_{\ell, X_1 \cap X_2}$. To see this observe first that if we compute the pullback of the sequence along the open immersion $j: X \setminus (X_1 \cap X_2) \hookrightarrow X$, then $\mathcal{F}$ vanishes and the morphism on the left becomes an isomorphism. In particular, $\mathcal{F}$ is supported on $X_1 \cap X_2$. To compute the pullback along the closed immersion $i: X_1 \cap X_2 \hookrightarrow X$ we may apply proper base change, since f is proper. But now the restriction of f to $X_0 := X_1 \cap X_2$ is simply the trivial double cover $X_0 \sqcup X_0 \twoheadrightarrow X_0$, so that $i^* \mathcal{F} \cong \mathbb{Q}_{\ell, X_0}$. This proves the claim because $\mathcal{F} \cong i_* i^* \mathcal{F}$, as $\mathcal{F}$ is supported on $X_1 \cap X_2$.

By an inductive application of the long exact cohomology sequences to sequences like (1), it suffices to prove the theorem for schemes X that are geometrically integral and separated algebraic over K . In this case, the proof follows by resolving X by a smooth hypercovering $X_\bullet$ see [6, p. 51], [8, 6.2.5] and the proof of [4, Thm. 6.3.2]. Since the hypercovering yields a spectral sequence that computes the cohomology of X in terms of the cohomologies of the smooth X_i , and this for all ℓ , and since only those X_i with $i \leq 2 \dim X$, contribute to X , the induction step is complete since we have reduced the case of arbitrary X to lower dimensions and to smooth X_i . $\square$

7 Proof of the main theorem

In this final section, we study the family $(\rho_{\ell, X})_{\ell \in L}$ in the particular case where K is absolutely finitely generated with the constant field k . We shall establish properties $\mathcal{R}(k)$ and if $\text{char}(k) > 0$ also property $\mathcal{S}(k)$. This will use Deligne's purity results from Weil I (cf. [7, Thm. 1.6]) as well as the global Langlands conjecture for function field proved by Lafforgue (cf. [26]).

Let k be a finite field and U a smooth k -variety. For every closed point $u \in U$ let $k(u)$ be the (finite) residue field of u , and let $D(u) \subset \pi_1(U)$ be the corresponding decomposition group (defined only up to conjugation). Denote by $\text{Fr}_u \in D(u)$ the preimage under the canonical isomorphism $D(u) \cong \text{Gal}(k(u))$ of the map $\sigma_u: x \mapsto x^{|k(u)|}$, so that within $\pi_1(U)$, the automorphism Fr_u is also defined only up to conjugation.

Proposition 7.1. *Suppose K is a finitely generated field of characteristic $p \geq 0$. Let k be the field of constants of K . Let X/K be a smooth projective scheme.*

- a) *There exists a finite separable extension K'/K and smooth k -variety U'/k with function field K' such that for each $j \geq 0$ and every $\ell \in \mathbb{L} \setminus \{p\}$, the representation $\rho_{\ell, X}^{(j)}|_{\text{Gal}(K')}$ of $\text{Gal}(K')$ is unramified along U' .*
- b) *If $p > 0$, then the family of representations $(\rho_{X, \ell}^{(j)}|_{\text{Gal}(K')})_{\ell \in \mathbb{L} \setminus \{p\}}$ is strictly compatible and pure of weight j , that is: For every closed point $u' \in U'$ the characteristic polynomial $p_{u'}(T)$ of $\rho_{\ell, X}^{(j)}(\text{Fr}_{u'})$ has integral coefficients, is independent of $\ell \in \mathbb{L} \setminus \{p\}$, and the reciprocal roots of $p_{u'}(T)$ all have absolute value $|k(u')|^{j/2}$.*

Proof. Note that k is perfect. There exists a finite separable extension E/K such that X_E splits up into a disjoint union of geometrically connected smooth projective E -varieties. Thus, after

replacing K by a finite separable extension, we can assume that X/K is geometrically connected. Let $(S_1, \dots, S_r)$ be a separating transcendence base of K/k . Identify $k(S_1, \dots, S_r)$ with the function field of $\mathbb{A}_r$ and let S be the normalization of $\mathbb{A}_r$ in K . Then S is a normal k -variety with function field K . There exists an alteration $S' \rightarrow S$ such that S'/k is a smooth variety and the function field K' of S' is a finite separable extension of K (cf. [6, Thm. 4.1, Remark 4.2]). Let $\mathfrak{A}$ be the set of all affine open subschemes of S' . Then $\mathrm{Spec}(K') = \varinjlim_{U' \in \mathfrak{A}} U'$. There exists

$U' \in \mathfrak{A}$ and a projective U' -scheme $f : \mathcal{X}' \rightarrow U'$ of such that $\mathcal{X}' \times_{U'} \mathrm{Spec}(K') = X_{K'}$. (cf. [17, 8.8.2] and [17, 8.10.5(v) and (xiii)]). By the theorem of generic smoothness, after shrinking U' and $\mathcal{X}'$, we can assume that f is smooth. Furthermore, after removing the support of $f_*(\mathcal{O}_{\mathcal{X}'})$ from U and shrinking $\mathcal{X}'$ accordingly, we can assume that f has geometrically connected fibres.

Let $j \geq 0$ and $u' \in U'$. Define $k' := k(u')$ and $X_{u'} := \mathcal{X}' \times_{U'} \mathrm{Spec}(k')$. Then for every $\ell \in \mathbb{L} \setminus \{p\}$ the étale sheaf $R^j f_* \mathbb{Z}_\ell$ is lisse and compatible with any base change (cf. [29, VI.2., VI.4]). Thus $\rho_{\ell, X}^{(q)}|_{\mathrm{Gal}(K')}$ factors through $\pi_1(U')$ and a) holds true. Furthermore it follows that $H^j(X_{\tilde{K}}, \mathbb{Q}_\ell)$ can be identified with $H^j(X_{u', \tilde{k}'}, \mathbb{Q}_\ell)$ in a way compatible with the Galois actions. Assume $p > 0$. Part b) then follows applying Deligne's theorem on the Weil conjectures (cf. [7, Thm. 1.6]) to $X_{u'}$. $\square$

Lemma 7.2. *Let the notation be as in the previous proposition and suppose we are in the situation of part (b), so that k is a finite field. Fix $q \in \mathbb{Z}$ and denote by n the dimension of $H^q(X_{\tilde{K}}, \mathbb{Q}_\ell)$ which is independent of $\ell \neq p$. Then the following hold:*

- (a) *For any smooth curve C over k and morphism $\varphi : C \rightarrow U'$, there exist irreducible cuspidal automorphic representations $(\pi_{C,i})_{i=1, \dots, m_\varphi}$ of $\mathrm{GL}_{n_i}(\mathbb{A}_{k(C)})$ such that $\sum_i n_i = n$ and for all ℓ the representations $(\varphi_* \circ \rho_{X, \ell}^{(q)})^{\mathrm{ss}}$ for $\varphi_* : \pi_1(C) \rightarrow \pi_1(U')$ agrees with the ℓ -adic representation $\oplus_i \rho_{\ell, \pi_{C,i}}$ attached to $\pi_{C,i}$ via the global Langlands correspondence in [26, p. 2].*
- (b) *If for some prime $\ell_0 \geq 3$ there is a $\mathbb{Z}_{\ell_0}$ -lattice L of the $\mathbb{Q}_{\ell_0}$ representation space underlying $\rho_{\ell_0, X, q}$ that is stabilized by $\mathrm{Gal}(K')$ and such that $\mathrm{Gal}(K')$ acts trivially on $L/\ell_0 L$, then in (a) for all $\varphi : C \rightarrow U'$ the representations $\pi_{C,i}$ are semistable⁴ at all places of $P(C) \setminus C$. In particular, for all $\ell \neq p$, the representation $\rho_{\ell, X}^{(q)}$ satisfies $\mathcal{S}(k)$.*

Proof. Let C be a smooth curve over k and $\varphi : C \rightarrow U'$ a morphism over k . By Proposition 7.1, the family of representations $(\varphi_* \circ \rho_{X, \ell}^{(q)})^{\mathrm{ss}}$, where $\ell \neq p$, is pure of weight j , semisimple and strictly compatible as a representation of $\pi_1(C)$. By the main theorem of [26, p. 2], each $(\varphi_* \circ \rho_{X, \ell}^{(q)})^{\mathrm{ss}}$ being pure and semisimple gives rise to a list of irreducible cuspidal automorphic representations

⁴We call an automorphic representation π of GL_n *semistable* at a place v if under the bijective local Langlands correspondence between local representations and Frobenius semi-simplified Weil-Deligne parameters, the Weil-Deligne parameter of π_v has underlying Weil-representation which is unramified. This definition is for instance used in [33, § 2]. In *loc. cit.* on page 8 it is also pointed out that by [3], this notion of semistability is equivalent to that of the automorphic representation having an Iwahori fixed vector.

via the global Langlands correspondence.⁵ By the strict compatibility and the bijectivity of the correspondence on simple objects, this list is the same for all ℓ (up to permutation). This proves (a).

Let now ℓ_0 be as in (b). For each $\varphi: C \rightarrow U'$ as in (a) consider the representation $(\varphi_* \circ \rho_{X, \ell_0}^{(q)})$ as an action on the lattice L , that is trivial modulo $\ell_0 L$. Any filtration of $L \otimes_{\mathbb{Z}_{\ell_0}} \mathbb{Q}_{\ell_0}$ preserved by the action of $\text{Gal}(k(C))$ induces a filtration of L . Denote by L_C the induced lattice for $(\varphi_* \circ \rho_{X, \ell_0}^{(q)})^{\text{ss}}$. Then it follows that the induced action of $\text{Gal}(k(C))$ on $L_C/\ell_0 L_C$ is trivial. Since $\ell_0 > 2$, this implies that series of the logarithm converges on the image of $(\varphi_* \circ \rho_{X, \ell_0}^{(q)})^{\text{ss}}$. This image being pro- ℓ , a standard argument (cf. [38, Cor. 4.2.2]) shows that the Weil-Deligne representation at any place of $P(C)$ attached to $(\varphi_* \circ \rho_{X, \ell_0}^{(q)})^{\text{ss}}$ has trivial underlying Weil-representation. But then by the compatibility of the global and local Langlands correspondence (cf. [26, Thme. VII.3, Cor. VII.5]), this means that all $\pi_{C,i}$ are semistable at the places in ∂C (they are unramified at all other places). We now apply the same argument to all $\ell \neq p$ in reverse, to deduce that all ramification of $(\varphi_* \circ \rho_{X, \ell}^{(q)})^{\text{ss}}$ is ℓ -tame for all $\ell \neq p$. The point simply is that in a family of Galois representations arising from a set of automorphic forms, the Weil-Deligne representation at a place of $P(C)$ is independent of $\ell \neq p$.

Finally from Proposition 4.5(b), which is a variant of a result of Kerz-Schmidt-Wiesend, we deduce that for all $\ell \neq p$, the representation $\rho_{\ell, X}^{(q)}$ is divisor ℓ -tame. Combined with Proposition 7.1(b), this establishes $\mathcal{S}(k)$. $\square$

Remark 7.3. We would like to point out the parallel between the proof of $\mathcal{S}(k)$ in part (b) of the previous lemma and in Example 5.2. In the proof of (b) we selected a prime ℓ_0 and enlarged K so that its image would act trivially on $L/\ell_0 L$ via ρ_{ℓ_0} for a $\text{Gal}(K)$ -stable lattice L . Then we could use the uniformity provided by automorphic representations (after restricting the ρ_{ℓ} to any curve) to deduce from this that all ramification was semistable in a sense.

In Example 5.2 we selected a prime ℓ_0 and enlarged K so that it contains $K(A[\ell_0])$. This requirement is equivalent to asking that $\text{Gal}(K)$ acts trivially on the quotient $T_{\ell_0}(A)/\ell_0 T_{\ell_0}(A)$ for the lattice $T_{\ell_0}(A)$ from the Tate-module. Then we used the uniformity provided by the Néron model $\mathcal{N}$ of the abelian variety A over any discrete valuation ring in K to deduce $\mathcal{S}(k)$. The semistability of $\mathcal{N}$ follows from the condition at the single prime ℓ_0 – in the analogous way that the semistability of automorphic representations was deduced from a single prime ℓ_0 .

Corollary 7.4. *Suppose K is absolutely finitely generated and of characteristic $p > 0$. Suppose X/K is smooth projective. Then $(\rho_{\ell, X})_{\ell \in \mathbb{L} \setminus \{p\}}$ satisfies $\mathcal{S}(k)$.*

Proof. Fix a prime $\ell_0 \geq 3$. Choose a lattice L underlying the representation $\rho_{\ell_0, X}$. Replace K by a finite extension such that $\text{Gal}(K)$ acts trivial on L/ℓ_0 via $\rho_{\ell_0, X}$. Apply now (b) of the previous lemma to deduce the corollary. $\square$

⁵Lafforgue in [26] describes only a correspondence between $\mathbb{Q}_{\ell}$ -sheaves of weight 0, i.e., Galois representations to GL_n whose determinant has finite image, and automorphic representations with finite order central character. A general statement is given in [9, § 0 and 0.1]

Theorem 7.5. *Let k' be a field of characteristic $p \geq 0$ and let K'/k' be a finitely generated extension. Let $L = \mathbb{L} \setminus \{p\}$. Let X'/k' be a separated algebraic scheme. Then there exists a finite extension E'/K' with the following properties:*

- (i) *For every $\ell \in L$ the group $\rho_{\ell, X'}(\text{Gal}(\tilde{k}'E'))$ lies in $\Sigma_\ell(2^n)$ and is generated by its ℓ -Sylow subgroups.*
- (ii) *The family $(\rho_{\ell, X'}|_{\text{Gal}(\tilde{k}'E')})_{\ell \in L \setminus \{2,3\}}$ is independent and $(\rho_{\ell, X'}|_{\text{Gal}(\tilde{k}'K)})_{\ell \in L}$ is almost independent.*

Proof. It suffices to establish conditions (a) and (b) of Corollary 6.7. For this, let K be absolutely finitely generated, that k is its field of constants and that X/K is smooth projective. In this case we have proven that $\mathcal{R}(k)$ holds in Proposition 7.1 and that $\mathcal{S}(k)$ holds for k of positive characteristic in Corollary 7.4. This verifies condition (a) of Corollary 7.4.

For condition (b) define n to be the dimensions of $\rho_{\ell, X}$. Since X/K is smooth projective, this definition is independent of the chosen ℓ . Then all images $\rho_{\ell, X}(\text{Gal}(K))$ are n -bounded at ℓ . Hence by Theorem 3.4, for each ℓ there exists a short exact sequence

$$1 \rightarrow M_\ell \rightarrow \rho_{\ell, X}(\text{Gal}(K)) \rightarrow H_\ell \rightarrow 1$$

with M_ℓ in $\Sigma_\ell(2^n)$ and H_ℓ in $\text{Jor}_\ell(J'(n))$, for the constant $J'(n)$ from Theorem 3.4. Consider the induced representations τ_ℓ defined as the composite $\text{Gal}(K) \xrightarrow{\rho_{\ell, X}} \rho_{\ell, X}(\text{Gal}(K)) \rightarrow H_\ell$. Since $\rho_{\ell, X}$ satisfies $\mathcal{R}(k)$ if $\text{char}(k) = 0$ and $\mathcal{S}(k)$ if $\text{char}(k) > 0$, the hypotheses of Proposition 4.7 are met if we replace $\text{Gal}(K)$ by $\pi_1(U')$ with U' from Proposition 7.1. Hence there exists a finite extension K' of K such that $\tau_\ell(\text{Gal}(K'\tilde{k}))$ is trivial for all $\ell \neq p$. But this shows that $\rho_{\ell, X}(\text{Gal}(K'\tilde{k}))$ lies in $\Sigma_\ell(2^n)$ for all $\ell \neq p$, proving condition (b) of Corollary 7.4 and completes the proof. $\square$

Remark 7.6. Under the hypothesis of the above theorem, Corollary 6.7 also tells us that $\mathcal{R}(k)$ holds for $(\rho_{\ell, X})_{\ell \in \mathbb{L}}$ if $\text{char}(k) = 0$ and that $\mathcal{S}(k)$ holds for $(\rho_{\ell, X})_{\ell \in \mathbb{L}}$ if $\text{char}(k) = p > 0$.

References

- [1] Emil Artin. The orders of the classical simple groups. *Comm. Pure and Applied Math.*, 8:455–472, 1955.
- [2] Michael Artin, Alexander Grothendieck, and Jean-Luis Verdier. *Séminaire de Géométrie Algébrique 4 - Théorie des topos et cohomologie étale des schémas*. Springer **LNLM** 269, 270, 305, 1972.
- [3] Joseph Bernstein and Andrey V. Zelevinsky. Induced representations of reductive p -adic groups. I. *Ann. Sci. École Norm. Sup. (4)*, 10:441–472, 1977.

- [4] Pierre Berthelot. Altération de variétés algébriques. *Sém. Bourbaki*, no. 815: 273–311, 1995-96.
- [5] Anna Cadoret, Akio Tamagawa. A refined Jordan-Nori theorem and applications to representations of étale fundamental groups. Preprint.
- [6] Johan Aise de Jong. Smoothness, semi-stability and alterations. *Publ. Math. IHES*, 83:51–93, 1996.
- [7] Pierre Deligne. La conjecture de Weil I. *Publ. Math. IHES*, 43:273–307, 1974.
- [8] Pierre Deligne. Théorie de Hodge III. *Publ. Math. I.H.E.S.*, 44:5–77, 1975.
- [9] Pierre Deligne. Finitude de l’extension de $\mathbb{Q}$ engendrée par des traces de Frobenius, en caractéristique finie. Volume dedicated to the memory of I. M. Gelfand. *Moscow Math. J.* 12, 2012.
- [10] Michel Demazure and Alexander Grothendieck. *Séminaire de Géométrie Algébrique 3 - Schemas en groups. Tome II*. Springer **LNM 152**, 1962/64.
- [11] Anna Devic and Richard Pink. Adelic openness for Drinfeld modules in special characteristic. *J. Number Theory* 132:1583–1625, 2012.
- [12] Mark Dickinson. On the modularity of certain 2-adic Galois representations. *Duke Math. J.*, 109:319–383, 2001.
- [13] Michael D. Fried and Moshe Jarden. *Field arithmetic. 2nd revised and enlarged ed.* Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge 11. Berlin: Springer. xxii, 780 p., 2005.
- [14] Wojciech Gajda and Sebastian Petersen. Independence of ℓ -adic Galois representations over function fields. *To appear in Compositio Mathematica*.
- [15] Alexander Grothendieck. Éléments de géométrie algébrique I (rédigé avec la coopération de Jean Dieudonné): I. Le langage des schémas. *Publ. Math. IHES*, (4):5–228, 1960.
- [16] Alexander Grothendieck. Éléments de géométrie algébrique (rédigé avec la coopération de Jean Dieudonné): II. Étude globale élémentaire de quelques classes de morphismes. *Publ. Math. IHES*, (8):5–222, 1961.
- [17] Alexander Grothendieck. Éléments de géométrie algébrique (rédigé avec la coopération de Jean Dieudonné): IV. Étude locale des schémas et des morphismes des schémas, Troisième partie. (28):5–255, 1966.
- [18] Alexander Grothendieck. *Séminaire de Géométrie Algébrique 1 - Revêtements étales et groupe fondamental*. Springer **LNM 224**, 1971.
- [19] Alexander Grothendieck. *Séminaire de Géométrie Algébrique 7 - Groupes de monodromy en géométrie algébrique*. Springer **LNM 288**, 1972.

- [20] Jin-Icho Igusa. *Fibre systems of jacobian varieties (III. Fibre systems of elliptic curves)*, Amer. J. Math. 81:453–476, 1959.
- [21] Luc Illusie. *Constructibilité générique et uniformité en ℓ* . Preprint.
- [22] Nicholas Katz and Serge Lang. Finiteness theorems in geometric class field theory. *Enseign. Math.*, 27(3-4):285–319, 1981.
- [23] Nicholas Katz and Gerard Laumon. Transformation de Fourier et majoration de sommes exponentielles. *Publ. Math. IHES*, 62:361–418, 1986; Erratum: *Publ. Math. IHES*, 69:244, 1989.
- [24] Mortiz Kerz and Alexander Schmidt. On different notions of tameness in arithmetic geometry. Preprint, [arXiv:0807.0979](https://arxiv.org/abs/0807.0979).
- [25] Wolfgang Kimmerle, Richard Lyons, Robert Sandling, and David Teague. Composition factors from the group ring and Artin’s theorem on orders of simple groups. *Proceedings LMS*, 60:89–122, 1990.
- [26] Laurent Lafforgue. Chtoucas de Drinfeld et correspondance de Langlands. *Inv. Math.* (147):1–241, 2002.
- [27] Michael Larsen and Richard Pink. Finite subgroups of algebraic groups. *Journal of the American Mathematical Society.*, (24), no. 4:1105–1158, 2011.
- [28] Werner Lütkebohmert. On compactification of schemes. *Manuscripta Math.* (80): 95–111, 1993.
- [29] James Milne. *Étale Cohomology*. Princeton University Press, 1980.
- [30] Gunter Malle and Donna Testerman. *Linear algebraic groups and finite groups of Lie type*. Cambridge studies in advanced mathematics 133, 2011.
- [31] Madhav Nori. On subgroups of $GL_n(\mathbb{F}_p)$. *Invent. math.*, (88):257–275, 1987.
- [32] Fabrice Orgogozo. Sur les propriétés d’uniformité des images en cohomologie étale. Preprint.
- [33] Conjeeveram Srirangachari Rajan. On Langlands functoriality – reduction to the semistable case. Algebraic groups and arithmetic, 199–219, TIFR, Mumbai, 2004.
- [34] Jean-Pierre Serre. Lettre à Ken Ribet du 7/3/1986. *Collected Papers IV*.
- [35] Jean-Pierre Serre. Une critère d’indépendance pour une famille de représentations ℓ -adiques. Preprint available at www.arxiv.org: 1006.2442.
- [36] Jean-Pierre Serre. propriétés conjecturales des groupes de Galois motiviques et des représentations ℓ -adiques. *Proc. Symp. Pure Math.*, (55):377–400, 1994.

- [37] Michel Vaquié. Valuations, in Resolution of singularities (Obergurgl, 1997). *Progr. Math.* (181): 539–590, Birkhäuser, Basel, 2000.
- [38] John Tate. Number Theoretic Background. *Proceedings Symp. Pure Math.* (33):3–26, 1979.

GEBHARD BÖCKLE
 COMPUTATIONAL ARITHMETIC GEOMETRY
 IWR (INTERDISCIPLINARY CENTER FOR SCIENTIFIC COMPUTING)
 UNIVERSITY OF HEIDELBERG
 IM NEUENHEIMER FELD 368
 69120 HEIDELBERG, GERMANY
 E-mail address: sekretariat.boeckle@iwr.uni-heidelberg.de

WOJCIECH GAJDA
 FACULTY OF MATHEMATICS AND COMPUTER SCIENCE
 ADAM MICKIEWICZ UNIVERSITY
 UMULTOWSKA 87
 61614 POZNAŃ, POLAND
 E-mail adress: gajda@amu.edu.pl

SEBASTIAN PETERSEN
 FB 10 - MATHEMATIK UND NATURWISSENSCHAFTEN
 UNIVERSITÄT KASSEL
 HEINRICH-PLETT-STR. 40
 34132 KASSEL, GERMANY
 E-mail adress: petersen@mathematik.uni-kassel.de